

GLP-1 Agonist/SGLT2 Hold Times

***** Failure to hold these medications could cause your procedure/surgery to be cancelled *****

Section 1: GLP-1A

(Glucagon-Like Peptid-1 Receptor Agonist)

If you are taking one of these medications or a compounded form, please hold them as indicated on this list. These medications slow the emptying of your stomach, which can put you at risk for aspiration when fully sedated.

Generic Name	Brand Name	Frequency	Days to <u>HOLD</u> before Surgery
Liraglutide	Victoza	Daily	1
Liraglutide	Saxenda	Daily	1
Semaglutide	Rybelsus	Daily	1
Exenatide (IR)	Byetta	Twice Daily	1
Semaglutide	Ozempic	Weekly	7
Semaglutide	Wegovy	Weekly	7
Dulaglutide	Trulicity	Weekly	7
Exenatide (ER)	Bydureon BCise	Weekly	7
Tirzepatide	Mounjaro	Weekly	7

Section 2: SGLT2

(Sodium-Glucose Cotransporter 2 Inhibitors)

Please stop any of the following medications 4 days prior to surgery to prevent potentially life-threatening postoperative ketoacidosis.

Generic Name	Brand Name	Days to <u>HOLD</u> before Surgery
Dapagliflozin	Farxiga	4
Empagliflozin	Jardiance	4
Ertugliflozin	Steglatro	4
Canagliflozin	Invokana	4
Bexagliflozin	Brenzavvy	4

Date & Version #	Change Summary
09/13/2024 Ver. 1	Original
10/10/2025 Ver 2	Added SGLT2